

REVIEW

Open Access



# Fluid bolus therapy in pediatric sepsis: a narrative review

Julian San Geroteo<sup>1\*</sup> , Michaël Levy<sup>1,2</sup>, Julien Gotchac<sup>3</sup>, Olivier Brissaud<sup>3</sup> and Stéphane Dager<sup>1,2</sup>

## Abstract

Leading cause of death in children under five, pediatric sepsis remains a significant global health threat. The 2020 Surviving Sepsis Campaign guidelines revised the management of septic shock and sepsis-associated organ dysfunction in children. In addition to empiric broad-spectrum antibiotics, fluid bolus therapy is one of the cornerstones of management, due to theoretical improvement of cardiac output, oxygen delivery and organ perfusion. Despite a very low level of evidence, the possible benefit of balanced crystalloids in sepsis resuscitation has led to discussion on their position as the ideal fluid. However, the latest adult data are not consistent with this, and the debate is still ongoing in pediatrics. We provide here the current state of knowledge on fluid bolus therapy in pediatric sepsis with emphasis on balanced crystalloids.

**Keywords:** Pediatric sepsis, Children, Fluid bolus therapy, Balanced crystalloids, Surviving sepsis campaign

## Introduction

Pediatric sepsis incidence and mortality are, respectively, 25.2 million and 3.4 million worldwide in 2017. More than half of sepsis cases and a third of sepsis-related deaths worldwide occur in children particularly in those under 5 years of age [1]. A recent large European prospective study of 2844 children with life-threatening infections reported a median age of 28 months in the cohort of 1229 pediatric sepsis [2]. The global case-fatality rate of pediatric sepsis is about 13% subjected to geographic variations [3, 4]. Sepsis accounts approximately for 3–8% of Pediatric Intensive Care Unit (PICU) admissions [5, 6]. Microbiological confirmation was obtained in only half of children, bacterial in about 45% of cases [2]. Two thirds of sepsis occurs in children with chronic disease who have higher in-hospital mortality [7].

To date, pediatric sepsis definition reference remains that of the 2005 consensus conference, which classified sepsis as infection in presence of systemic inflammatory

response syndrome [8]. The lack of sensitivity and specificity of these historical criteria [9, 10] led to a change in the adult definition of sepsis as a “life-threatening organ dysfunction caused by a dysregulated host response to infection” [11]. However, this concept is not yet validated in pediatrics [12], and therefore, the latest guidelines of the Surviving Sepsis Campaign refer to children with infection and cardiovascular (i.e. “septic shock”) or non-cardiovascular organ dysfunction (i.e. “other sepsis-associated organ dysfunction”) [13].

Sepsis bundles include an early fluid resuscitation, the obtention of a quick vascular access, collection of blood cultures, rapid initiation of broad-spectrum antibiotics, the measure of lactate and early administration of vasoactive agents if shock persists [13]. Completion of sepsis bundles lead in decreasing sepsis-related mortality [14].

Although the use of normal saline remains predominant in pediatric sepsis [15, 16], balanced crystalloids are now suggested as the first-line fluid in the latest guidelines of the Surviving Sepsis Campaign [13] and the European Resuscitation Council [17]. However, the ideal fluid debate is still ongoing [18]. In this narrative review, we aim to provide an overview of the current state of knowledge regarding fluid bolus therapy in pediatric

\*Correspondence: [julian.san-geroteo@aphp.fr](mailto:julian.san-geroteo@aphp.fr)

<sup>1</sup> Pediatric Intensive Care Unit, Robert-Debré University Hospital, Paris, France  
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

sepsis, with emphasis on available data regarding balanced crystalloids.

### Fluid bolus therapy indications

World Health Organization (WHO) recommendation in the 1990s to perform aggressive fluid bolus therapy reduced tenfold the mortality of pediatric sepsis [19]. Fluid bolus therapy improves cardiac output, oxygen delivery, and organ perfusion in sepsis [20], although such a response is variable and unpredictable [21, 22]. Fluid bolus therapy indications in pediatric sepsis have recently been revised in the Surviving Sepsis Campaign [13].

In healthcare systems with access to intensive care, fluid bolus therapy is recommended in case of abnormal perfusion or hypotension for age. Clinical signs of poor tissue perfusion are tachycardia, altered mental status, decreased urine output (below 1 ml/kg/h) and abnormalities in capillary refill time (flash < 1 s or prolonged > 3 s), skin (mottled, cool or flushed, ruddy) and pulses (weak or bounding) [23]. Despite being a standard of care in high income countries, the evidence for fluid bolus therapy in pediatric sepsis is limited in these settings [24] and is based on historical studies of weak methodology [25].

In healthcare systems without access to intensive care, fluid bolus therapy is only recommended in case of hypotension, defined as low systolic blood pressure (i.e., < 50 mmHg before 1 year, < 60 mm Hg between 1 and 5 years and < 70 mm Hg after 5 years) or all three WHO shock criteria (i.e., cold extremities, capillary refill time greater than 3 s and weak or fast pulse) [13]. This major distinction regarding fluid bolus therapy indications in low- and middle-income countries, where intensive care is not routinely accessible, comes mainly from the FEAST (*Fluid Expansion As Supportive Therapy*) study, the only randomized clinical trial (RCT) to date that has evaluated fluid bolus therapy in pediatric sepsis. Maitland et al. randomized 3141 sub-Saharan African children aged 2 months to 12 years with sepsis and abnormal perfusion without severe hypotension to, either fluid bolus therapy (20 mL/kg NaCl 0.9% or 5% albumin) or no bolus with maintenance fluid only. After exclusion of severe malnutrition or gastroenteritis, mortality at 48 h and 1 month was higher with fluid bolus therapy. However, the findings of the FEAST should be interpreted cautiously [26, 27]. Indeed, the WHO shock criteria were absent in one third of children and lack of PICU availability worsens the outcome of fluid-related complications. Moreover, a transfusion threshold of 5 g/dL, not reached in one third of cases, and the diagnosis of malaria in more than half of the cases limit the external validity of this study in high resource settings, especially since fluid bolus

therapy could be dangerous in these situations because of hemodilution [28].

Finally, in both healthcare systems, fluid bolus therapy should be discontinued as soon as signs of fluid overload develop, such as pulmonary oedema or new or worsening hepatomegaly. Moreover, cumulative positive fluid balance is associated with poor outcomes in children [29, 30], especially with sepsis [31–33].

In summary, according to the Surviving Sepsis Campaign, fluid bolus therapy indications in pediatric sepsis are limited to hypotension, regardless of health systems, and abnormal perfusion (only in settings with availability of intensive care) [13]. Once the decision for volume expansion has been made, one of the key elements that may affect children's outcome, is the fluid choice for sepsis initial resuscitation.

### Resuscitation fluid choice: crystalloid solutions

#### Rationale

Composed of water and electrolytes, crystalloid solutions (Table 1) are divided in normal saline (or NaCl 0.9% and "physiological" saline) and balanced crystalloids. Normal saline is a high-chloride solution with 154 mmol/L of both Na<sup>+</sup> and Cl<sup>-</sup>, main cation and anion of the body, respectively. Balanced crystalloids are low-chloride solutions, closer to the electrolyte composition of plasma, where Cl<sup>-</sup> anions are replaced by buffers (lactate, acetate, gluconate, or malate), quickly excreted, or metabolized into bicarbonate. According to Stewart approach, acid–base balance is mainly determined by the strong ion difference, i.e. the difference between all fully dissociated cations ([Na<sup>+</sup>] + [K<sup>+</sup>] + [Ca<sup>2+</sup>] + [Mg<sup>2+</sup>]) and anions ([Cl<sup>-</sup>] + [lactate] + [SO<sub>4</sub><sup>2-</sup>]) [34]. Physiologically, plasma strong ion difference is about 40 mmol/L (total strong cation > strong anion concentrations). Below 24 mmol/L, the lower the strong ion difference, the more dissociation of H<sub>2</sub>O into H<sup>+</sup> and OH<sup>-</sup> occurs, leading to hyperchloremic metabolic acidosis. Because of its zero SID ([Na<sup>+</sup>] = [Cl<sup>-</sup>]), the use of normal saline drives to a lower bicarbonate concentration, a shift of H<sup>+</sup> and K<sup>+</sup> out of cells for electroneutrality and hyperchloremic metabolic acidosis. Balanced crystalloids have strong ion difference > 24 mmol/L with a neutral or alkalinizing action on pH, due to lower Cl<sup>-</sup> concentration than Na<sup>+</sup> [35, 36].

Hyperchloremia decreases tubular chloride reabsorption, resulting in arteriolar vasoconstriction, and therefore, a decrease in glomerular filtration rate and urine output [37]. Apart from the kidney, chlorine-rich fluids are associated with increased inflammation, coagulopathy, macrocirculatory and microcirculatory haemodynamic alterations [38].

**Table 1** Composition of commonly used crystalloids

|             | Human plasma | Normal saline | Hartmann's solution | Ringer's lactate | Ringer's acetate | Plasma-Lyte | Sterofundin |
|-------------|--------------|---------------|---------------------|------------------|------------------|-------------|-------------|
| Sodium      | 140±5        | 154           | 131                 | 130              | 130              | 140–141     | 145         |
| Chloride    | 102±8        | 154           | 111                 | 109              | 112              | 98          | 127         |
| Potassium   | 4.5±1        | 0             | 5                   | 4                | 5                | 5           | 4           |
| Calcium     | 2.4±0.2      | 0             | 2                   | 1.4              | 1                | 0           | 2.5         |
| Magnesium   | 0.9±0.1      | 0             | 0                   | 0                | 1                | 1.5         | 1           |
| Bicarbonate | 28±4         | 0             | 0                   | 0                | 0                | 0           | 0           |
| Lactate     | <2           | 0             | 29                  | 28               | 0                | 0           | 0           |
| Gluconate   | 0            | 0             | 0                   | 0                | 0                | 23          | 0           |
| Acetate     | 0            | 0             | 0                   | 0                | 27               | 27          | 24          |
| Malate      | 0            | 0             | 0                   | 0                | 0                | 0           | 5           |
| Osmolarity  | 285±10       | 308           | 278                 | 273              | 276              | 295         | 309         |
| pH          | 7.4±0.02     | 4.5–7.0       | 5.0–7.0             | 6.0–7.5          | 6.0–8.0          | 6.5–8.0     | 5.1–5.9     |
| SID         | 40±2         | 0             | 27                  | 27               | 25               | 50          | 27          |
| Na/Cl ratio | 1.21–1.54    | 1             | 1.18                | 1.19             | 1.16             | 1.43        | 1.14        |

All variables are expressed in mmol/L except for osmolarity (mosm/L) and pH. SID strong ion difference

Hyperchloremia and hyperchloremic metabolic acidosis are common in PICU, occurring in 95% and 39% of children, respectively [39]. Hyperchloremia or even moderate increase in  $[Cl^-] \geq 5$  mmol/L has been associated with poor outcomes in children in several retrospective studies [40–43].

The Surviving Sepsis Campaign recommends, with very low quality of evidence, balanced crystalloids rather than normal saline, as the first-line for fluid bolus therapy in pediatric sepsis [13]. The latest guidelines of the European Resuscitation Council have recently extended this recommendation to all childhood circulatory failures [17].

In children without sepsis, a recent meta-analysis of Lehr et al. (13 studies, 11,848 children) comparing balanced crystalloids (Ringer's Lactate in 8/13 studies) and normal saline showed a significant improvement regarding pH and serum bicarbonate level (mean increase of 0.03 and 1.6 mmol/L, respectively) with balanced crystalloids in the pooled analysis of 3 RCTs with acute gastroenteritis. No differences were found in metabolic acidosis, hyperchloremia, acute kidney injury, need for mechanical ventilation, vasopressor, or renal replacement therapy, as well as PICU or hospital length of stay and mortality. Interpretation of this work is difficult because of the multiple analyses performed on studies with small sample sizes, highly heterogeneous patients and endpoints, two of which were at high risk of bias [44].

In pediatric sepsis, a recent Indian RCT of 708 children with septic shock found a reduction in the incidence of acute kidney injury, need for renal

replacement therapy and hyperchloremia but not modification on survival with Plasma-Lyte [45]. Most of the other existing data are from two large American retrospective studies with equivocal results. Emrath et al. found a reduction in mortality, acute kidney injury and need for vasopressors with balanced crystalloids at 72 h in 7000 children [46], whereas Weiss et al. showed no difference on acute kidney injury or mortality between normal saline and balanced crystalloids in 4234 pediatric sepsis but increase hospital length of stay with balanced crystalloids. Nevertheless, in this last study, the severity of children who received balanced crystalloids (older and statistically greater age, crystalloid volume, cardiovascular and respiratory disease, steroids, albumin, and transfusion), and the small number of children (459/4234) who received “only balanced crystalloids” suggests that this was a second-line treatment and makes the comparison difficult. In addition, the median volume of crystalloid of 24 ml/kg, may have diminished the power of the study [47]. Finally, a recent observational study of 99 pediatric sepsis found a decrease in acute kidney injury, need for renal replacement therapy and hospital length of stay with balanced crystalloids [48]. Thus, the debate is not over and the results of an ongoing international RCT with anticipated enrollment of 8,800 children with septic shock are eagerly awaited [49], especially as two recent large RCTs in adults ICUs did not report improved survival with balanced crystalloids [50, 51].

### Clinical practice

In clinical practice, as Ringer's Lactate is slightly hypotonic (Table 1), others balanced crystalloids or normal saline should be used instead in patients at risk of increased intracranial pressure (e.g., traumatic brain injury, diabetic ketoacidosis). Normal saline should be preferred in hypovolemic hyponatremia or hypochloremic contraction with metabolic alkalosis due to its higher  $[\text{Na}^+]$  and  $[\text{Cl}^-]$  content. In all other situations, balanced crystalloids are suggested as first-line crystalloids, especially in pediatric sepsis [13, 17]. They should be used with caution in cases of anuria, although hyperkalemia does not contraindicate their use in critically ill children [52]. Indeed, balanced crystalloids do not generate higher kalemia, because their  $[\text{K}^+]$  content is rapidly diluted in the extracellular fluid and the  $[\text{K}^+]$  shifting out of cells induced by normal saline-related hyperchloremic metabolic acidosis is much greater [53, 54]. Balanced crystalloids must be infused on a different route from red blood cell transfusions or Ceftriaxone as they contain calcium. Regarding side effects, balanced crystalloids should not be considered equivalent one to another and have their own adverse effects [55]. For example, lactate buffered balanced crystalloids may induce small rises in serum lactate concentration in patient with severe liver failure [56]. Similarly, high acetate concentrations have been associated with hypotension and direct myocardial toxicity [57]. Finally, balanced crystalloids do not generate additional costs in comparison with normal saline [58].

### Comparison between crystalloids and colloids

Although fluid bolus therapy with albumin seems to be non-inferior to crystalloids, its high cost, lack of availability, blood-borne infectious risk and high-chloride content do not allow its routine use to be recommended for initial fluid bolus therapy [13, 17], except in specific situations, such as severe dengue because of extensive capillary leaks [59]. Finally, other colloids should be abandoned in pediatric sepsis and other childhood circulatory failures [13, 17]. Starches use have been completely ruled out by the European Medicines Agency because of the risk of anaphylaxis, coagulopathy and acute kidney injury in adults [60] and gelatin have shown no benefit in the only RCT carried out in pediatric sepsis [61].

In summary, balanced crystalloids, which are closer to the composition of plasma with a lower chlorine content than normal saline, their theoretical reduction in the risk of hyperchloremic metabolic acidosis and subsequent acute kidney injury, inflammation, and coagulopathy are now suggested as the first-line choice for fluid bolus

therapy in pediatric sepsis and other childhood circulatory failures.

### Fluid volume bolus and duration

Fluid bolus therapy is still recommended by the Surviving Sepsis Campaign in case of abnormal perfusion or hypotension for age in healthcare systems with access to intensive care [13], although related to excess mortality in a recent meta-analysis of 19 studies involving 9,321 cases of pediatric sepsis. This is likely due to that Yue et al. included mainly studies from low- to middle-income countries, such as India ( $n=6$ ), Kenya ( $n=5$ ) or Vietnam ( $n=3$ ). Moreover, this result was not consistent in the subgroup of general shock patients after exclusion of malaria case [62].

### Volume

The Surviving Sepsis Campaign recommends fluid boluses of 10–20 ml/kg of ideal body weight, up to 40–60 ml/kg over the first hour of management [13]. The European Resuscitation Council have recently extended this recommendation to all childhood circulatory failures, preferably with boluses of 10 ml/kg [17]. These smaller boluses comparing with previous guidelines [63, 64], are argued by the need for more frequent and faster clinical reassessment as well as the early use of vasoactive agents. However, it does not limit the total amount of fluid bolus therapy to be given. Even smaller volumes of 5–10 ml/kg may be beneficial to optimize preload in given pathologies such as pre-existing heart disease at risk of acute pulmonary edema, under the condition of even more frequent clinical reassessments [65]. This reconsideration of the 20 ml/kg paradigm is supported by a multicenter pilot study on 75 children with sepsis assigned to bolus of 10 ml/kg or 20 ml/kg after an initial bolus of 20 ml/kg for both groups, where no differences were found in mortality, hospital length of stay, PICU transfer, receipt of mechanical ventilation or inotropes and duration of mechanical ventilation [66]. The target of 40–60 ml/kg over the first hour of management through repeated boluses was associated with increased survival [25] and decreased occurrence of hepatomegaly [67] in pediatric historical studies. Finally, in adults, there are no prospective intervention comparing different volumes for initial resuscitation in sepsis [68] and one retrospective study showed that failure to receive 30 mL/kg of fluid bolus therapy within 3 h of sepsis onset was associated with increased mortality, delayed hypotension and ICU stay [69].

### Duration

The Surviving Sepsis Campaign recommends fluid boluses of 10–20 ml/kg of ideal body weight up to 40–60 ml/kg

over the first hour of management in pediatric sepsis and no longer mentions the “5–10 min” duration as previously for fluid bolus therapy. We argue that this short duration should be maintained to achieve the above-mentioned goal of 1-h 40–60 ml/kg fluid bolus therapy even if no supportive studies in children clearly confirm the benefit in using 5 to 10 min rather than 10 to 20 min for fluid bolus therapy. A recent large multicentric Brazilian RCT with more than 10,000 critically ill adults found no difference in mortality after bolus at a rate of 999 ml/h versus 333 ml/h [70].

## Conclusion

Pediatric sepsis remains a significant global health threat, although revision to the reference 2005 children definition is still lacking. Management of pediatric sepsis have been updated by the 2020 Surviving Sepsis Campaign, whose compliance lead in decreasing sepsis-related mortality. These international guidelines notably suggest balanced crystalloids as the first-line solution for initial fluid bolus therapy. However, there is a very low level of evidence supporting this recommendation to date. Thus, the ideal fluid debate is not over, and the results of an ongoing large pediatric randomized clinical trial are eagerly awaited.

## Author contributions

JSG contributed to the initial draft of the manuscript and created table. All authors read and approved the final manuscript.

## Funding

Pediatric Intensive Care Unit of Robert-Debré University Hospital.

## Availability of data and materials

Not applicable.

## Declarations

## Ethics approval and consent to participate

Not applicable.

## Consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## Author details

<sup>1</sup>Pediatric Intensive Care Unit, Robert-Debré University Hospital, Paris, France.

<sup>2</sup>Université Paris Cité, Paris, France. <sup>3</sup>Neonatal and Pediatric Intensive Care Unit, Children's University Hospital, Bordeaux, France.

Received: 6 September 2022 Accepted: 4 November 2022

Published online: 12 November 2022

## References

- Rudd KE, Johnson SC, Agesa KM, Shackelford KA, Tsoi D, Kievlan DR, et al. Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the Global Burden of Disease Study. *Lancet*. 2020;395:200–11.
- Martinón-Torres F, Salas A, Rivero-Calle I, Cebey-López M, Pardo-Seco J, Herberg JA, et al. Life-threatening infections in children in Europe (the

- EUCLIDS Project): a prospective cohort study. *Lancet Child Adolesc Health*. 2018;2:404–14.
- Tan B, Wong JJ-M, Sultana R, Koh JC, Jit M, Mok YH, et al. Global case-fatality rates in pediatric severe sepsis and septic shock: a systematic review and meta-analysis. *JAMA Pediatr*. 2019;173:352–62.
- Fleischmann-Struzek C, Goldfarb DM, Schlattmann P, Schlapbach LJ, Reinhart K, Kissoon N. The global burden of paediatric and neonatal sepsis: a systematic review. *Lancet Resp Med*. 2018;6:223–30.
- Weiss SL, Fitzgerald JC, Pappachan J, Wheeler D, Jaramillo-Bustamante JC, Salloo A, et al. Global epidemiology of pediatric severe sepsis: the sepsis prevalence, outcomes, and therapies study. *Am J Respir Crit Care Med*. 2015;191:1147–57.
- Schlapbach LJ, Straney L, Alexander J, MacLaren G, Festa M, Schibler A, et al. Mortality related to invasive infections, sepsis, and septic shock in critically ill children in Australia and New Zealand, 2002–13: a multicentre retrospective cohort study. *Lancet Infect Dis*. 2015;15:46–54.
- Prout AJ, Talisa VB, Carcillo JA, Mayr FB, Angus DC, Seymour CW, et al. Children with chronic disease bear the highest burden of pediatric sepsis. *J Pediatr*. 2018;199:194–199.e1.
- Goldstein B, Giroir B, Randolph A, International Consensus Conference on Pediatric Sepsis. International pediatric sepsis consensus conference: definitions for sepsis and organ dysfunction in pediatrics. *Pediatr Crit Care Med*. 2005;6:2–8.
- Weiss SL, Fitzgerald JC, Maffei FA, Kane JM, Rodriguez-Nunez A, Hsing DD, et al. Discordant identification of pediatric severe sepsis by research and clinical definitions in the SPROUT international point prevalence study. *Crit Care*. 2015;19:325.
- Scott HF, Deakynne SJ, Woods JM, Bajaj L. The prevalence and diagnostic utility of systemic inflammatory response syndrome vital signs in a pediatric emergency department. *Acad Emerg Med*. 2015;22:381–9.
- Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 2016;315:801–10.
- Morin L, Hall M, de Souza D, Guoping L, Jabornisky R, Shime N, et al. The current and future state of pediatric sepsis definitions: an international survey. *Pediatrics*. 2022;149: e2021052565.
- Weiss SL, Peters MJ, Alhazzani W, Agus MSD, Flori HR, Inwald DP, et al. Surviving sepsis campaign international guidelines for the management of septic shock and sepsis-associated organ dysfunction in children. *Intensive Care Med*. 2020;46:10–67.
- Rodrigues-Santos G, de Magalhães-Barbosa MC, Raymundo CE, Lima-Setta F, da Cunha AJLA, Prata-Barbosa A. Improvement of 1st-hour bundle compliance and sepsis mortality in pediatrics after the implementation of the surviving sepsis campaign guidelines. *J Pediatr (Rio J)*. 2021;97:459–67.
- Gelbart B, Schlapbach L, Ganeshalingham A, Ganu S, Erickson S, Oberender F, et al. Fluid bolus therapy in critically ill children: a survey of practice among paediatric intensive care doctors in Australia and New Zealand. *Crit Care Resusc*. 2018;20:131–8.
- Samransamruajkit R, Wong JJM, Smathakane C, Anantasi N, Sunkonkit K, Ong J, et al. Pediatric severe sepsis and shock in three Asian countries: a retrospective study of outcomes in nine PICUs. *Pediatr Crit Care Med*. 2021;1:1.
- Van de Voorde P, Turner NM, Djakow J, de Lucas N, Martinez-Mejias A, Biarent D, et al. European resuscitation council guidelines 2021: paediatric life support. *Resuscitation*. 2021;161:327–87.
- Fuhrman DY, Kellum JA. Hyperchloremic IV solutions: have we seen enough?...or “Still Good Medicine?” *Pediatr Crit Care Med*. 2018;19:171–2.
- Carcillo JA, Tasker RC. Fluid resuscitation of hypovolemic shock: acute medicine's great triumph for children. *Intensive Care Med*. 2006;32:958–61.
- Long E, Duke T. Fluid resuscitation therapy for paediatric sepsis. *J Paediatr Child Health*. 2016;52:141–6.
- Ranjit S, Natraj R, Kissoon N, Thiagarajan R, Ramakrishnan B, Monge Garcia MI. Variability in the Hemodynamic Response to Fluid Bolus in Pediatric Septic Shock. *Pediatr Crit Care Med*. 2021;
- Long E, Babl FE, Oakley E, Sheridan B, Duke T. Pediatric Research in Emergency Departments International Collaborative (PREDICT). Cardiac index changes with fluid bolus therapy in children with sepsis-an observational study. *Pediatr Crit Care Med*. 2018;19:513–8.

23. Long E, Babl F, Dalziel S, Dalton S, Etheridge C, Duke T, et al. Fluid resuscitation for paediatric sepsis: a survey of senior emergency physicians in Australia and New Zealand. *Emerg Med Australas*. 2015;27:245–50.
24. Gelbart B. Fluid bolus therapy in pediatric sepsis: current knowledge and future direction. *Front Pediatr*. 2018;6:308.
25. Carcillo JA, Davis AL, Zaritsky A. Role of early fluid resuscitation in pediatric septic shock. *JAMA*. 1991;266:1242–5.
26. Hilton AK, Bellomo R. A critique of fluid bolus resuscitation in severe sepsis. *Crit Care*. 2012;16:302.
27. Myburgh J, Finfer S. Causes of death after fluid bolus resuscitation: new insights from FEAST. *BMC Med*. 2013;11:67.
28. Duke T. What the African fluid-bolus trial means. *Lancet*. 2011;378:1685–7.
29. Alobaidi R, Basu RK, DeCaen A, Joffe AR, Lequier L, Pannu N, et al. Fluid accumulation in critically ill children. *Crit Care Med*. 2020;48:1034–41.
30. Alobaidi R, Morgan C, Basu RK, Stenson E, Featherstone R, Majumdar SR, et al. Association between fluid balance and outcomes in critically ill children: a systematic review and meta-analysis. *JAMA Pediatr*. 2018;172:257–68.
31. Abulebda K, Cvijanovich NZ, Thomas NJ, Allen GL, Anas N, Bigham MT, et al. Post-ICU admission fluid balance and pediatric septic shock outcomes: a risk-stratified analysis. *Crit Care Med*. 2014;42:397–403.
32. Bhaskar P, Dhar AV, Thompson M, Quigley R, Modem V. Early fluid accumulation in children with shock and ICU mortality: a matched case-control study. *Intensive Care Med*. 2015;41:1445–53.
33. Chen J, Li X, Bai Z, Fang F, Hua J, Li Y, et al. Association of Fluid Accumulation with Clinical Outcomes in Critically Ill Children with Severe Sepsis. *PLoS ONE*. 2016;11: e0160093.
34. Kishen R, Honoré PM, Jacobs R, Joannes-Boyau O, De Waele E, De Regt J, et al. Facing acid-base disorders in the third millennium - the Stewart approach revisited. *Int J Nephrol Renovasc Dis*. 2014;7:209–17.
35. Morgan TJ, Venkatesh B, Hall J. Crystalloid strong ion difference determines metabolic acid-base change during acute normovolaemic haemodilution. *Intensive Care Med*. 2004;30:1432–7.
36. Li H, Sun S, Yap JQ, Chen J, Qian Q. 0.9% saline is neither normal nor physiological. *J Zhejiang Univ Sci B*. 2016;17:181–7.
37. Lobo DN, Awad S. Should chloride-rich crystalloids remain the mainstay of fluid resuscitation to prevent “pre-renal” acute kidney injury?: con. *Kidney Int*. 2014;86:1096–105.
38. Semler MW, Kellum JA. Balanced Crystalloid Solutions. *Am J Respir Crit Care Med*. 2018;199:952–60.
39. Bulfon AF, Alomani HL, Anton N, Comrie BT, Rochweg B, Stef SA, et al. Intravenous fluid prescription practices in critically ill children: a shift in focus from natremia to chloremia? *J Pediatr Intensive Care*. 2019;8:218–25.
40. Barhight MF, Lusk J, Brinton J, Stidham T, Soranno DE, Faubel S, et al. Hyperchloremia is independently associated with mortality in critically ill children who ultimately require continuous renal replacement therapy. *Pediatr Nephrol*. 2018;33:1079–85.
41. Suetrong B, Pisitsak C, Boyd JH, Russell JA, Walley KR. Hyperchloremia and moderate increase in serum chloride are associated with acute kidney injury in severe sepsis and septic shock patients. *Crit Care*. 2016;20:315.
42. Barhight MF, Brinton J, Stidham T, Soranno DE, Faubel S, Griffin BR, et al. Increase in chloride from baseline is independently associated with mortality in critically ill children. *Intensive Care Med*. 2018;44:2183–91.
43. Stenson EK, Cvijanovich NZ, Anas N, Allen GL, Thomas NJ, Bigham MT, et al. Hyperchloremia is associated with complicated course and mortality in pediatric patients with septic shock. *Pediatr Crit Care Med*. 2018;19:155–60.
44. Lehr AR, Rached S, Barrowman N, Tsampalieros A, Parker M, McIntyre L, et al. Balanced versus unbalanced fluid in critically ill children: systematic review and meta-analysis. *Pediatr Crit Care Med*. 2022;1:1.
45. Sankar J, Jayashree M, Mahadevan S, et al. Balanced crystalloids versus saline for initial fluid resuscitation in children with septic shock. *Crit Care Med*. 2021;49:17–17.
46. Emrath ET, Fortenberry JD, Travers C, McCracken CE, Hebbar KB. Resuscitation with balanced fluids is associated with improved survival in pediatric severe sepsis\*. *Crit Care Med*. 2017;45:1177–83.
47. Weiss SL, Keele L, Balamuth F, Vendetti N, Ross R, Fitzgerald JC, et al. Crystalloid fluid choice and clinical outcomes in pediatric sepsis: a matched retrospective cohort study. *J Pediatr*. 2017;182:304–310.e10.
48. Fernández-Sarmiento J, Alcalá-Lozano C, Barrera PA, Erazo Vargas SC, Gómez Cortes LB, Reyes CM. Association between unbalanced solutions and acute kidney injury during fluid resuscitation in children with sepsis. *J Intensive Care Med*. 2022;37:625–32.
49. Weiss SL, Balamuth F, Long E, Thompson GC, Hayes KL, Katcoff H, et al. PRagMatic Pediatric Trial of Balanced vs nOrmal Saline FLUID in Sepsis: study protocol for the PRoMPT BOLUS randomized interventional trial. *Trials*. 2021;22:776.
50. Zampieri FG, Machado FR, Biondi RS, Freitas FGR, Veiga VC, Figueiredo RC, et al. Effect of intravenous fluid treatment with a balanced solution vs 0.9% saline solution on mortality in critically ill patients: the BaSICS randomized clinical trial. *JAMA*. 2021;326:818–29.
51. Finfer S, Micallef S, Hammond N, Navarra L, Bellomo R, Billot L, et al. Balanced multielectrolyte solution versus saline in critically ill adults. *N Engl J Med*. 2022;386:815–26.
52. Barhight MF, Nelson D, Moran T, Christiano J, Sanchez-Pinto LN. Association between the use of balanced fluids and outcomes in critically ill children: a before and after study. *Crit Care*. 2021;25:266.
53. Santi M, Lava SAG, Camozzi P, Giannini O, Milani GP, Simonetti GD, et al. The great fluid debate: saline or so-called “balanced” salt solutions? *Ital J Pediatr*. 2015;41:47.
54. Hayes W. Ab-normal saline in abnormal kidney function: risks and alternatives. *Pediatr Nephrol*. 2019;34:1191–9.
55. Curran JD, Major P, Tang K, Bagshaw SM, Dionne JC, Menon K, et al. Comparison of balanced crystalloid solutions: a systematic review and meta-analysis of randomized controlled trials. *Crit Care Explor*. 2021;3: e0398.
56. Shin W-J, Kim Y-K, Bang J-Y, Cho S-K, Han S-M, Hwang G-S. Lactate and liver function tests after living donor right hepatectomy: a comparison of solutions with and without lactate. *Acta Anaesthesiol Scand*. 2011;55:558–64.
57. Morgan TJ. The ideal crystalloid - what is “balanced”? *Curr Opin Crit Care*. 2013;19:299–307.
58. Taylor C, Yang L, Finfer S, Machado FR, YouZhong A, Billot L, et al. An international comparison of the cost of fluid resuscitation therapies. *Aust Crit Care*. 2021;34:23–32.
59. Ranjit S, Ramanathan G, Ramakrishnan B, Kissoon N. Targeted interventions in critically ill children with severe dengue. *Indian J Crit Care Med*. 2018;22:154–61.
60. European Medicines Agency. Hydroxyethyl-starch solutions for infusion to be suspended – CMDh endorses PRAC recommendation. 2018. <https://www.ema.europa.eu/en/news/hydroxyethyl-starch-solutions-infusion-be-suspended-cmdh-endorses-prac-recommendation>
61. Upadhyay M, Singhi S, Murlidharan J, Kaur N, Majumdar S. Randomized evaluation of fluid resuscitation with crystalloid (saline) and colloid (polymer from degraded gelatin in saline) in pediatric septic shock. *Indian Pediatr*. 2005;42:223–31.
62. Yue J, Zheng R, Wei H, Li J, Wu J, Wang P, et al. Childhood Mortality after fluid bolus with septic or severe infection shock: a systematic review and meta-analysis. *Shock*. 2021;56:158–66.
63. Davis AL, Carcillo JA, Aneja RK, Deymann AJ, Lin JC, Nguyen TC, et al. The American College of critical care medicine clinical practice parameters for hemodynamic support of pediatric and neonatal septic shock: executive summary. *Pediatr Crit Care Med*. 2017;18:884–90.
64. Dellinger RP, Levy MM, Carlet JM, Bion J, Parker MM, Jaeschke R, et al. Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2008. *Crit Care Med*. 2008;36:296–327.
65. Weiss SL, Nicolson SC, Naim MY. Clinical update in pediatric sepsis: focus on children with pre-existing heart disease. *J Cardiothorac Vasc Anesth*. 2020;34:1324–32.
66. Inwald DP, Canter R, Woolfall K, Mouncey P, Zenasni Z, O’Hara C, et al. Restricted fluid bolus volume in early septic shock: results of the Fluids in Shock pilot trial. *Arch Dis Child*. 2019;104:426–31.
67. Santhanam I, Sangareddi S, Venkataraman S, Kissoon N, Thiruvengadamudayan V, Kasthuri RK. A prospective randomized controlled study of two fluid regimens in the initial management of septic shock in the emergency department. *Pediatr Emerg Care*. 2008;24:647–55.

68. Evans L, Rhodes A, Alhazzani W, Antonelli M, Coopersmith CM, French C, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021. *Crit Care Med.* 2021;49: e1063.
69. Kuttab HI, Lykins JD, Hughes MD, Wroblewski K, Keast EP, Kukoyi O, et al. Evaluation and predictors of fluid resuscitation in patients with severe sepsis and septic shock. *Crit Care Med.* 2019;47:1582–90.
70. Zampieri FG, Machado FR, Biondi RS, Freitas FGR, Veiga VC, Figueiredo RC, et al. Effect of slower vs faster intravenous fluid bolus rates on mortality in critically ill patients: the basics randomized clinical trial. *JAMA.* 2021;326:830–8.

### **Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

