

CORRECTION

Open Access



Correction: Extracellular vesicles derived from endothelial cells modulate macrophage phenotype in vitro

Zhizhen He^{1†}, Johannes Greven^{1,4†}, Yulong Shi^{1,2*}, Kang Qin¹, Qun Zhao¹, Xing Zhang¹, Eva Miriam Buhl³, Jörg Eschweiler¹, Frank Hildebrand¹ and Elizabeth Rosado Balmayor^{1,4*}

Correction: European Journal of Medical Research (2023) 28:506

<https://doi.org/10.1186/s40001-023-01427-6>

In the original publication of the article, the Fig. 3 was duplicated as Fig. 1 inadvertently. The corrected Fig. 1 is given below. The original [1] article has been corrected.

Published online: 11 December 2023

[†]Zhizhen He and Johannes Greven have contributed equally to this work.

The original article can be found online at <https://doi.org/10.1186/s40001-023-01427-6>.

*Correspondence:

Yulong Shi
yulong.shi@whu.edu.cn
Elizabeth Rosado Balmayor
erosadobalma@ukaachen.de

¹ Department of Orthopedics, Trauma and Reconstructive Surgery, RWTH Aachen University Hospital, Pauwelsstraße 30, 52074 Aachen, Germany

² Division of Joint Surgery and Sports Medicine, Department of Orthopedic Surgery, Zhongnan Hospital of Wuhan University, Wuhan 430071, China

³ Electron Microscopy Facility, Institute for Pathology, RWTH Aachen University Hospital, Pauwelsstraße 30, 52074 Aachen, Germany

⁴ Experimental Orthopaedics and Trauma Surgery, RWTH Aachen University Hospital, Pauwelsstraße 30, 52074 Aachen, Germany



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 1 Characterization of isolated E-EVs. **A** TEM images of a representative sample of the isolated ELPS-EVs and ENor-EVs. Micrographs revealed a circular and double-membrane structure characteristic of EVs. **B** NTA determinations of size (nm) and concentration (particles/ml) of isolated ELPS-EVs and ENor-EVs. **C** Identification of EVs protein markers CD63 and CD31

Reference

1. He Z, Greven J, Shi Y, Qin K, Zhao Q, Zhang X, Buhl EM, Eschweiler J, Hildebrand F, Balmayor ER. Extracellular vesicles derived from endothelial cells modulate macrophage phenotype in vitro. *Eur J Med Res.* 2023;28:506. <https://doi.org/10.1186/s40001-023-01427-6>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.